

The circular seabird economy is critical for oceans, islands and people

Holly P. Jones^{1,2}✉, Jennifer Appoo³, Cassandra E. Benkwitt⁴, Stephanie B. Borrelle⁵, Ruth E. Dunn^{4,6}, Hannah E. Epstein⁷, Leilani A. Fowlke¹, Nick D. Holmes⁸, Laura-Li Jeannot⁴, Yadvinder Malhi^{9,10}, Lyndsay L. Rankin^{11,16}, Sebastian Steibl^{12,13}, Christy N. Wails¹⁴, David J. Will¹⁵ & Nicholas AJ Graham⁴

Abstract

Nearly a third of seabird species are at risk of extinction, rendering them among the most threatened bird taxa globally. The decline in seabird populations has major ramifications for their associated ecosystems. An expansive literature covers seabird impacts on their breeding islands, and growing research expands understanding of the circular seabird economy – cross-ecosystem connections of seabirds from oceans to islands, and back to nearshore marine ecosystems, mainly driven by nutrient transfer. Amid the twin biodiversity and climate crises, maintaining and restoring the circular seabird economy is one way to achieve large-scale, cross-ecosystem impacts with simple and time-tested conservation methods. This Review summarizes researchers' understanding of the circular seabird economy and outlines goals for future research. Historically, research has focused on terrestrial impacts, with emerging research explicitly studying the cross-ecosystem impacts of seabirds. We identify substantial knowledge gaps, with research needed to understand how the circular seabird economy changes with space, time, across biomes and with island size. Emerging analytic and monitoring tools will help to propel this research forward. Many coastal communities, especially Indigenous people, rely on seabirds for their livelihoods and cultural connections, and so taking a people-centred approach incorporating Indigenous knowledge and perspectives is critical moving forward.

Sections

Introduction

Marine nutrients from sea to land

Seabird impacts on islands

Seabird impacts on adjacent marine ecosystems

Supporting circular seabird economies

Summary and future directions

¹Department of Biological Sciences, Northern Illinois University, DeKalb, IL, USA. ²Institute for the Study of the Environment, Sustainability and Energy, Northern Illinois University, DeKalb, IL, USA. ³Blue Economy Research Institute, University of Seychelles, Mahé, Seychelles. ⁴Lancaster Environment Centre, Lancaster University, Lancaster, UK. ⁵BirdLife International, Pacific Secretariat, Suva, Fiji. ⁶Centre d'Ecologie Fonctionnelle et Evolutive, Montpellier, France. ⁷School of Life Sciences, University of Essex, Colchester, UK. ⁸California Chapter, The Nature Conservancy, Santa Cruz, CA, USA. ⁹Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, UK. ¹⁰Leverhulme Centre for Nature Recovery, University of Oxford, Oxford, UK. ¹¹Independent researcher, Davis, CA, USA. ¹²Naturalis Biodiversity Center, Leiden, The Netherlands. ¹³School of Biological Sciences, University of Auckland, Auckland, New Zealand. ¹⁴Department of Natural Resource Management, South Dakota State University, Brookings, SD, USA. ¹⁵Island Conservation, Santa Cruz, CA, USA. ¹⁶Present address: United States Geological Survey, Davis, CA, USA. ✉e-mail: hjones@niu.edu

Introduction

Seabirds are top predators in marine ecosystems, consuming comparable amounts of primarily fish and squid to the commercial fishing industry's annual take^{1–4}. Most seabirds spend the majority of their lives on the open ocean, but as obligate land breeders they come ashore to nest, predominantly on islands⁵. Seabirds are colonial and can nest in millions on a single island, although colonies typically range from hundreds to thousands of birds⁶. This movement of seabirds from oceans to land results in the movement of massive amounts of dietary-derived marine nutrients (carbon, nitrogen, phosphorus and calcium). These nutrients are deposited on islands via guano, eggshells, feathers and carcasses of chicks and adults, resulting in profound effects on island biodiversity and ecosystem functioning. Indeed, the impacts of seabirds on their breeding islands are so unique and consequential that they are often termed seabird islands⁷. However, the nutrient flow does not end on islands; the marine-derived nutrients eventually wash back into the ocean through runoff, groundwater seeps and, in the case of volatilized ammonia from guano, via wind. These nutrient inputs have important consequences for marine intertidal and nearshore environments, creating what has been previously termed the circular seabird economy⁸. The circular seabird economy is the process by which seabirds consume their prey in the ocean, transport nutrients to islands and then those nutrients are washed back out into the ocean, to be taken up by ocean organisms (Fig. 1). We argue this term provides a useful descriptor for the vast scale of seabird impacts in the many ecosystems that they link and helps those outside the disciplines of ecology and conservation to understand the importance of those links for the livelihoods of coastal communities dependent on seabird inputs.

Despite their importance for driving biodiversity on and around islands, and as top predators in oceans, seabirds are one of the most threatened taxa of birds, with nearly one-third listed as threatened, endangered or critically endangered⁹. Moreover, extinctions since European colonization have disproportionately occurred on islands¹⁰, which are the primary nesting location for seabirds. Given that climate change threatens their breeding islands and negatively affects their food availability^{9,11–13}, seabirds sit at the nexus of both the biodiversity and climate crises. Because seabird impacts span large scales and multiple ecosystems, conserving and restoring their populations is one relatively simple way to maintain beneficial large-scale cross-ecosystem impacts. To understand how best to use seabird conservation to address the biodiversity crisis, and given emerging science that an intact seabird island, in turn, could be an important local resilience factor for island ecosystems facing climate change, it is first important to understand how the circular seabird economy works.

In this Review, we summarize research around the circular seabird economy (Fig. 1), describing the flow of nutrients and their ecological impacts. We highlight the effects of the circular seabird economy on ecosystems adjacent to seabird islands and identify threats to this balanced system, as well as describing relevant conservation efforts. We identify knowledge gaps pertaining to the circular seabird economy, and thus areas for prospective research, and end with what can propel this area of research into the future.

Marine nutrients from sea to land

Seabirds comprise ~3.5% of all avifauna and are defined as birds that are reliant on marine habitats for elements of their life cycles¹⁴. Body size can vary greatly (from 20-g least storm-petrels, *Halocryptena*




Fig. 1 | The seabird circular economy. Seabirds extract marine nutrients from the sea and deposit them on islands. These nutrients are assimilated directly in the terrestrial ecosystem and back into nearshore marine systems via runoff.

This cycle is sensitive to biotic and abiotic factors that can vary with season, phenology and biogeography (Box 1).

microsoma, to 30–38-kg emperor penguins, *Aptenodytes forsteri*⁵) as well as foraging range (for example, <5 km in incubating Canadian roseate terns, *Sterna dougallii*, and >6,800 km in great shearwaters, *Ardenna gravis*, incubating eggs within the South Atlantic Ocean; Fig. 2). During their breeding seasons, seabirds are central place foragers, sometimes travelling vast distances from their nesting and roosting sites to feed, on trips that can last from hours to days, regularly crossing the terrestrial–marine boundary¹⁵.

Seabirds feed most prevalently on pelagic marine organisms, particularly krill, squid and fish². Globally, seabirds extract around 50–96 million tonnes of prey annually from marine ecosystems^{1–4}, securing energy for their basal metabolic processes, movement, growth, moult and reproduction. Seabird energy expenditure, and therefore their energetic requirements, increases with body mass¹⁶, and large-bodied species, such as albatrosses and penguins (particularly the genus *Aptenodytes*), require substantial quantities of prey to balance their energetic budgets (Fig. 2). Seabird energetics also vary spatiotemporally, contributing to seasonal pulses in subsequent nutrient subsidies, with energy expenditure generally increasing throughout the breeding season as adults care for their developing offspring¹⁶. Outside the breeding season, moult is an energetically costly period, during which many seabirds also return to land¹⁷, thereby depositing marine-derived nutrients that were assimilated into feather tissue from elsewhere in their ranges.

The breeding, life history and foraging behaviour of seabirds mean they function as biological pumps, consuming prey within their oceanic feeding grounds to meet their energetic requirements and then transferring large quantities of nutrients to their terrestrial roosting and breeding grounds^{18,19}. The concentrations and quantities of nutrients that seabirds deposit at their colonies are influenced by factors such as body size, rate of energetic expenditure and foraging ecology¹⁷, which can also vary by individual, sex, age and location²⁰ (Fig. 2). The protein-rich marine prey species of seabirds result in the transfer of two key elements: nitrogen and phosphorus²¹. Globally, seabirds transfer 3.8×10^3 Gg nitrogen and 0.63×10^3 Gg phosphorus per year from the ocean to land²². This nutrient flux is comparable in magnitude to other major global flows of nitrogen and phosphorus, such as non-anthropogenic oceanic inputs of nitrogen via groundwater and riverine inorganic phosphorus fluxes to the ocean²². Guano represents the main form of nutrient input at colonies, and once deposited, some of the nitrogen is emitted to the atmosphere as ammonia gas²³, particularly in hot and humid environments²⁴. In addition to nitrogen and phosphorus, seabird-derived nutrients contain a suite of elements such as zinc, copper, manganese, nickel, cadmium, lead and iron^{25–27}. Although most inputs from seabirds are biotic or part of natural geochemical cycling, anthropogenic pollution means that seabirds also transport plastic and other contaminants to their breeding colonies²⁸. Nutrient inputs via guano deposition can be substantial, with an estimated 591 Gg nitrogen, 99 Gg phosphorus, 39.3 Mg cadmium and 27.2 Mg lead being deposited annually at breeding colonies, demonstrating that seabirds are important components of numerous global biogeochemical cycles^{19,22}. The fate of deposited nutrients includes being taken up into island food webs via plants and consumers, especially detritivores^{29–31}, transported back into the marine environment through runoff^{28,29,30}, into groundwater^{31,32} and, in the case of volatilized ammonia, transported into nearby ecosystems via wind^{24,33,34}.

Seabird impacts on islands

The quantity of nitrogen and phosphorus deposition to seabird islands varies by seabird body size, energy requirements, population size, life




Fig. 2 | Seabird mass and foraging range. The relationship between mean bird mass (kg) and foraging range for breeding adult seabird species (km). Species are categorized by International Union for Conservation of Nature (IUCN) Red List conservation status (symbols), daily energy expenditure (kJ per day; point size) and taxonomic family (colour). Data on mean bird mass are from ref. 180. Data on mean seabird foraging ranges were collated for breeding adult seabirds where data exist within ref. 181 via Google Scholar searches ($n = 110$ seabird species). Where means were not available, medians were used. Daily energy expenditure was calculated via the Seabird FMR Calculator in ref. 16. See Supplementary information for further information and complete data sources. IUCN Red List categories: least concern (LC), near threatened (NT), vulnerable (VU), endangered (EN) and critically endangered (CR).

history and phenology. Typically, seabirds with larger populations, larger-bodied species and longer nesting periods (or year-round roosts) contribute larger amounts than others^{35–39}. Quantities taken up into island food webs and washed back into the ocean vary according to island abiotic factors such as island size, soil type, slope, precipitation regime and terrestrial nutrient status along the scale of oligotrophic to eutrophic^{8,40,41} (Box 1) and presence of other biotic sources of marine-derived nutrients such as beach wrack, pinnipeds and intertidal invertebrates⁴¹. For example, impacts in arid ecosystems are positively

Box 1 | Sources of variation in impacts for islands and marine links

Environmental and ecological factors affecting both seabird distribution and behaviour, as well as the input and residency time of seabird-derived nutrients in nearshore marine habitats, can influence the contribution, distribution and impact of nutrient flows across the land–sea interface.

Seabird ecology

Seabird-derived nutrient flows to marine habitats are ultimately governed by guano deposition that is linked with seabird diet and breeding behaviours. The nutrient composition and quantity of guano can be dictated by species-specific feeding and nesting preferences, seabird body size, colony size and density and breeding season¹²¹. Guano deposition on islands and in nearshore marine habitat can therefore be concentrated in areas in and around colonies^{55,120}, might be temporally or seasonally limited¹³⁸, and could vary with seabird diversity and community composition³⁷. Additionally, these natural nutrient flows can be disrupted by the presence of island invaders, whereby invasive vegetation restricts breeding habitat^{44,182} and invasive predators eat seabirds (reviewed elsewhere^{183–185}).

Island dynamics

The physical characteristics of islands, such as their total area, topography, elevation, slope and watershed size, further affect seabird nesting and roosting patterns and the cross-ecosystem flow of nutrients (Fig. 1). Given that nutrients can enter marine habitats through watershed or surface runoff and through percolation and submarine groundwater discharge^{186,187}, these island dynamics, coupled with variations in annual or seasonal precipitation, become key factors in determining the concentrations and patterns of nutrient distribution in nearshore marine ecosystems. For example, high islands typically have larger watersheds than low-lying islands or atolls, such that the primary vector for nutrient input into nearshore habitat can differ between runoff or stream outflow and submarine groundwater discharge, respectively¹⁸⁸.

Windscaapes

Global and local wind patterns have ubiquitous influence on both seabird behaviour and nearshore marine hydrodynamics that can

cause further variability in nutrient dynamics across the land–sea interface. Prevailing winds such as the easterly trade winds in the tropics, doldrums near the Equator and mid-latitude westerlies can affect large-scale migratory patterns and seabird distribution, as well as flight, foraging and breeding behaviours (reviewed elsewhere¹⁸⁹). High winds and storms might also disrupt guano deposition by creating inaccessible nesting grounds¹⁹⁰. Both prevailing winds and storms also increase wave exposure that more readily impacts windward compared with leeward sides of islands, perhaps increasing the presence of internal waves^{191,192} and decreasing the residency time of nutrients in nearshore waters¹⁹³.

Seascapes

Other physical features of the nearshore marine habitat can influence the dissipation of nutrients and their residency times, and ultimately the degree of impact, or size of impact zone, of seabird-derived nutrients on marine ecosystems. Differences have been observed in the bioaccumulation of seabird-derived nutrients into algal, coral or fish tissue among varying distances from shore^{29,112,122}, reef habitats and depth^{40,194} and at distance from active seabird colonies¹¹⁵. Biogeophysical modelling of seabird nutrient fluxes to marine systems has further identified variables such as seafloor curvature and water flow as crucial to local nutrient-delivery patterns¹⁹⁵.

Biomes

Each biome has unique seabird diversity, ecology, island dynamics, windscaapes and seascapes, driving marked differences in the circular seabird economy between biomes. Seabird-derived nutrient impacts tend to be stronger in the oligotrophic systems of polar, desert and tropical ecosystems versus relatively eutrophic temperate systems⁷¹⁹⁶. Precipitation is a driving force of variation in seabird-derived nutrient impacts, with higher precipitation yielding stronger seabird-derived nutrient impacts across different biomes^{40,197}. The degree of guano volatilization varies across biomes and will affect transfer to nearshore ecosystems; for example, in the hot, humid tropics, most guano quickly volatilizes into ammonia rather than flowing into ecosystems through soil deposition^{22,24,198}.

correlated with rainfall⁴², and impacts on temperate islands are stronger in the rainy season^{8,40}. Because the impacts of nitrogen and phosphorus deposition vary according to precipitation and nutrient context, they also vary by biome (Table 1 and Box 1), with stronger impacts seen in relatively nutrient-poor polar regions versus nutrient-rich temperate regions^{26,43}, and the relatively few studies available suggesting substantial impacts in nutrient-poor tropical systems^{37,44,45}.

Seabird-derived inputs can alleviate terrestrial nutrient limitation, enhancing primary productivity and carbon sequestration, although effects vary with vegetation types, climates and along longitudinal gradients. On tropical and subtropical islands, early successional vegetation can respond markedly to seabird influence⁴⁶, whereas mature, densely forested systems exhibit weaker responses^{47,48}. Effects on primary production are modulated by rainfall, with positive effects on plant cover and biomass under sufficient rainfall⁴⁹ but toxic effects of seabird guano prevailing under dry conditions, often lowering plant

cover⁵⁰. In nutrient-limited Arctic systems, seabirds can increase carbon stocks⁵¹, plant cover⁵² and plant biomass^{43,53}. Plant biomass increases coincide with compositional shifts towards plant species and traits adapted to uptake of seabird guano. Guano deposition can trigger transitions from woody to grass-dominated vegetation in tropical⁵⁴, temperate⁵⁵ and Arctic systems^{52,56}, often reducing species richness^{57,58}. Under high guano loads, monodominant vegetation stands can form on tropical atoll islands (such as *Pisonia* forests)^{59,60}, as well as in Arctic systems⁵⁶. Seabirds facilitate seed dispersal, actively driving compositional shifts in plant community composition^{61–63}. Elevated terrestrial plant biomass can change herbivory rates, although plant trait-related factors govern general response strengths and directions⁶⁴.

Increased carbon stocks and guano loads alter soil composition, nutrient content and functioning across biomes and longitudinal gradients. Nitrogen and phosphorus generally increase, whereas soil pH decreases^{49,51}. Increased microbial biomass can transition from nitrogen

Review article

to carbon limitation, thus accelerating leaf litter decomposition⁶⁵ and soil nutrient and carbon fluxes^{66,67}. These indirect effects of seabirds on island soils, via elevated nutrient inputs and terrestrial primary production, are often paired with direct soil impacts of seabirds, such as biopedturbation. Seabird burrowing can increase or decrease soil water content, depending on geography, vegetation cover and detritus accumulation^{57,58}. Trampling and burrowing can reduce seedling species richness and density^{68,69}, modify soil compaction⁵⁷ and accelerate erosion⁴⁹.

In general, seabirds are associated with greater animal species diversity and abundance on islands, and positive impacts for ecosystems^{26,70–72}. Invertebrate and lizard community composition and species richness are altered through changes in habitat and predator dynamics where seabirds are present. For example, on Gulf of California islands, seabirds promote increased invertebrate consumer abundance and increased abundance for their corresponding spider and lizard predators⁷³. Invertebrates and detritivores, which directly consume seabird tissues and guano, show the strongest abundance increases⁷⁴. Seabird impacts on terrestrial biodiversity are greatest on smaller islands, where seabirds are a unique or primary source of nutrients and where terrestrial ecosystems have otherwise limited internal nutrient stocks relative to a highly productive marine ecosystem⁷³.

Given that breeding seabirds affect the abundance of island biota, it should be no surprise that those impacts yield unique food web interactions. For example, increased diversity and abundance of detritivores, ectoparasites and scavengers that feed directly from seabird carcasses can yield more nutrient resources for their prey⁷⁵. Impacts on consumers are expected to vary with biomes such that nutrient-rich systems and/or drier biomes exhibit lower-magnitude impacts to food webs than nutrient-poor and/or wetter biomes (Box 1 and Table 1). In nutrient-rich temperate ecosystems, seabirds serve as prey for many native island species; for example, around 8% of the large endemic Phillip Island centipede's (*Cormocephalus coynei*) diet comprises black-winged petrel (*Pterodroma nigripennis*) chicks⁷⁶. Likewise, seabird burrows provide habitat for other species, especially reptiles such as tuatara (*Sphenodon punctatus*), lizards and snakes^{77–80}. In Arctic biomes, whole-ecosystem transformations occur when seabird-derived nutrients pulse through these otherwise nutrient-limited systems, the whole ecosystem transforms from tundra to grassland, with attendant impacts to island biota, as seen in the Aleutians^{43,53,56}. Native deer mice (*Peromyscus maniculatus*) depredate Scripps's murrelet (*Synthliboramphus scrippsi*) chicks⁸¹, and scores of native birds of prey and other seabirds consume nesting seabirds, eggs and chicks^{82–85}.

Seabird impacts on adjacent marine ecosystems

The influence of seabirds on the coastal ecosystems that fringe their islands is a more nascent field²⁶. Identified links between seabirds and intertidal community effects date back to the 1980s^{86,87}, but more substantive research has only emerged in the past decade^{40,88,89}. Linkages are typically close to islands, in shallow water ecosystems such as coral reefs, seagrasses, rocky shores and mangroves⁹⁰. Most links between seabirds on islands and nearshore coastal ecosystems are through the transfer of nutrients. Nutrients including nitrogen and phosphorus, and other elements found in seabird guano such as iron, enter coastal ecosystems via runoff, through groundwater discharges, and via gas exchange⁴¹. The influence of these nutrient exchanges for ecosystem functions, food web structures and biodiversity can be substantial. The strength of nutrient signals and uptake by marine organisms are

mediated by processes such as wave energy and oceanic nutrient influence^{40,91} (Box 1).

Seabird nutrient beneficiary ecosystems are often different from the marine ecosystems where the majority of seabirds are feeding and sourcing energy to transfer back to islands⁹². Indeed, for most seabirds, which feed some distance away from islands (Fig. 1), the circular seabird economy might not be a closed feedback loop, with the source marine energy coming from different spatial scales and ecosystems than the marine habitats that ultimately benefit. Alternatively, for seabirds that feed closer to shore, the circular seabird economy could function more like a closed feedback loop with seabird prey benefiting from locally seabird-derived nutrients. Notably, even pelagic feeding seabirds can affect forage prey density within 150 km from their breeding colonies⁹³. The contribution of fish from nearshore environments to the seabird diet can be quite minimal, despite these communities being the most affected by breeding seabird nutrients^{92,94}. However, seabird-derived nutrients can benefit nearshore habitats such as seagrass meadows, which serve as nurseries for younger age classes of seabird to forage fish⁹⁴, potentially closing the feedback loop. For islands in dry environments, with little potential for runoff, the conduit for nutrient transfer to the sea is via wind transporting volatilized ammonia^{19,21,95}. For seabirds breeding on inland islands far from the ocean, there is no nutrient transfer back to the ocean and thus no closed loop. Identifying what influences the extent to which the seabird circular economy is a closed loop is a key future target for this field.


Where seabirds are providing otherwise limiting nutrients, seabird inputs can enhance growth rates of individual marine organisms (Fig. 3). For example, guano addition stimulates primary production of phytoplankton from coastal and open ocean areas^{96–98} and macroalgae in the rocky intertidal zone⁹⁹. Primary producers from a range of marine ecosystems also have higher abundance, biomass and/or per cent cover near seabird colonies (such as polar phytoplankton, subtropical seagrasses and temperate intertidal macroalgae)^{86,96,100}. Although less is known about seabird nutrient effects on higher trophic levels (reviewed elsewhere^{101,102}), on coral reefs seabird nutrients increase coral growth^{103–105}, fish growth^{29,88,106} and fish biomass⁸⁸. However, effects on biomass and abundance can be variable both within and among trophic levels (Table 1 and Box 1), in part owing to species interactions such as competition and predation^{107,108}. For example, macroalgal cover has been shown to increase, decrease and remain similar across reefs with varying seabird nutrient inputs, probably owing to

Table 1 | Relative strength of terrestrial impacts of seabirds across biomes


Climate zone	Plant diversity	Plant biomass	Animal diversity	Animal biomass	Food web
Arctic	--	+++	?	?	?
Northern Hemisphere temperate	--	++	++	++	++
Tropical	--	+	+++	+	?
Southern Hemisphere temperate	--	++	++	++	++
Sub-Antarctic	--	+++	?	?	?

Impacts assessments are based on studies included in this Review, with strength of the impact denoted with positive (+) or negative (–) notation. Each impact varies by factors discussed in Box 1. Many remain understudied, as indicated by question marks (?).


a Seabird impacts on marine ecosystems



b Individual responses



c Food web interactions



d Biodiversity




Fig. 3 | Seabird impacts on adjacent marine ecosystems. a, Impacts on temperate (left) and tropical (right) ecosystems. **b,** Coral and reef fish growth rates are boosted by seabird nutrients. **c,** Seabird presence reshapes food web

interactions in a mangrove. **d,** Macroalgae diversity in a temperate intertidal system increases near seabird colonies. Data in part **b** are from refs. 88,103,104. Data in part **c** are from ref. 116. Data in part **d** are from ref. 33.

differences in available space and/or herbivory^{109–112}. Similarly, even within a single system seabird nutrients can have different effects across species and trophic levels, resulting in altered community structure rather than a uniform increase in abundance or biomass across all organisms^{30,107,113,114}.

Seabird nutrient inputs enrich nearshore coastal food webs across multiple trophic levels^{29,30,115–117} (Fig. 4). Like in terrestrial ecosystems, nutrient enrichment of primary producers in coastal and marine habitats, and their increase in biomass and abundance, can enhance herbivory rates. For example, in mangrove forests, seabird-derived nutrients boost mangrove productivity and improve leaf nutritional quality, leading to increased herbivory by folivorous insects¹¹⁸. On coral reefs, herbivorous parrotfish exhibit higher

grazing rates on algae enriched by seabird-derived nutrients⁸⁸. Seabird nutrient inputs also modify marine producer–consumer interactions by altering feeding pathways. In seabird-subsidized food webs, consumers can shift their resource use towards primary producers with higher concentrations of seabird-derived nitrogen. Such resource shifts have been observed across multiple ecosystems and taxa, including invertebrates in coastal ponds¹¹⁹, mangroves⁴⁵ and nearshore waters^{120,121}, as well as cryptobenthic fishes inhabiting shallow coral reefs¹⁰⁶. The hard coral *Pocillopora damicornis* adjusts its feeding strategy in response to seabird-derived nutrient availability, relying more on autotrophy when nutrient input is high, while shifting towards mixotrophy or heterotrophy when inputs are lower, depending on seasonal and site-specific

conditions¹²² (Box 1). Additionally, seabird nutrient inputs can alter interactions by influencing consumer behaviour. For example, on coral reefs near seabird islands, damselfish feeding on turf algae enriched with seabird-derived nutrients show reduced foraging effort and smaller foraging territories while exhibiting more aggressive territory defence, compared with those on reefs near islands without seabirds¹²³.

In addition to species growth and food web interactions, seabird nutrient subsidies can also affect coastal biodiversity (Fig. 3). These impacts can differ in direction and intensity, with both positive and negative relationships identifiable in biodiversity and ecosystem function comparisons (such as biomass and productivity) across terrestrial and aquatic systems¹²⁴. In coastal systems, guano enrichment can alter microbial¹²⁵ and phytoplankton¹¹⁴ community structure and growth. The addition of nitrogen-rich guano in laboratory experiments resulted in an overall biomass increase and shifting species dominance⁹⁸. Comparisons of natural seabird islands reveal slightly greater macroalgae species richness at high-density seabird islands³⁰ and seabird islands with the longest history of seabird presence relative to lower densities and shorter presence histories⁴⁰. However, nearshore coral reefs show decreasing macroalgal richness, evenness and diversity with increasing seabird-derived nutrients¹¹². Seagrass beds are more diverse with extensive coverage surrounding seabird islands compared with islands without seabirds¹⁰⁰. Seabird nutrient subsidies do not alter biodiversity and ecosystem functioning relationships in coral reef fishes, but instead enhance ecosystem functions directly, and have marginal positive effects on biodiversity⁸⁹. Biodiversity and ecosystem functioning relationships

are highly dependent on the local ecosystem, trophic interactions and spatial scale^{126–128}. Seabird nutrient subsidies could also influence biodiversity metrics differently based on biome (polar, temperate and tropical).

Supporting circular seabird economies

The circular seabird economy faces multiple ongoing and interacting threats. However, there are innovative methods to help to restore seabirds and their linkages to islands and oceans and ensure that the circular seabird economy is resilient to ongoing threats. Global climate change is a particularly pressing threat to the circular seabird economy, because it affects seabirds directly through sea level rise and increased storm frequency on low-lying breeding islands and indirectly through modulating seabird prey availability. Seabirds themselves might confer adaptive capacity to the terrestrial and marine ecosystems they connect, which will be increasingly important in an increasingly uncertain future.

Current threats to future resilience

Past and on-going anthropogenic activities at sea and on land have led to seabird extinctions, extirpations from breeding islands and reduced abundance on land and sea¹²⁹, thus interrupting the seabird circular economy for many land–ocean systems. Comprehensive syntheses of threats to seabirds on land and at sea have classified threat timing, scope and severity by species and identified where knowledge gaps exist^{9,129}. Invasive predators remain the key threat to nearly half of seabird species worldwide⁹. Seabirds killed in fisheries, or bycatch, are the second most important threat to albatrosses and petrels. Other threats include overfishing of prey species, climate change altering prey




Fig. 4 | Seabird restoration methods. Predator exclusion fencing prevents invasive mammalian predators from accessing the newly established seabird colonies. Plantings of native plant species can provide habitat for seabirds to roost. Use of decoys can attract seabirds to suitable translocation sites. Auditory

social attraction play-back systems with long-range speakers, a solar panel and battery can also be used for attracting seabirds to translocation sites. Finally, translocation interventions can support struggling populations, including tube feeding of chicks or establishment of interspecific surrogacy programmes.

distributions¹³⁰, direct harvesting of chicks and eggs⁹, light pollution¹³¹ and plastic pollution/ingestion^{132–134}. The threats that seabirds face are increasing and intensifying, and novel threats are emerging, such as highly pathogenic diseases, pollution and offshore wind farm developments¹³⁵. Although studies are increasingly investigating the cumulative and synergistic impacts of these multiple and dynamic threats to seabirds on land and at sea, understanding their complex interplay remains a considerable challenge¹¹.

Seabird restoration reconnects oceans and islands

Despite threats to seabirds, there are tools to ensure their future resilience by protecting existing colonies, increasing existing populations and establishing new colonies. Conserving existing colonies, establishing protected areas in and around those colonies and investing in biosecurity are common sense approaches. Removing invasive predators, either through eradication or predator-proof fencing, is often the first step to seabird recovery (Fig. 4) and has been shown to re-establish the circular seabird economy in many instances^{29,136}. Once predators are removed, one way to kickstart seabird colonies is to restore native island vegetation so that seabirds have suitable nest-building materials, structural support for their burrows and nests and the improved ability to avoid avian predators^{137,138} (Fig. 4).

Beyond restoring vegetation, by far the most common ways to increase extant populations or establish new colonies are through direct social attraction and/or translocation^{95,139,140} (Fig. 4). Seabird social attraction uses visual, auditory, olfactory or a combination of these cues to entice target seabird species to (re)colonize and breed at a restoration site; this method has been used in over 800 restoration projects of at least 138 seabird species⁹⁵. Many seabirds depend on social cues, such as the presence of other seabirds to select a nesting location^{141–145}, so social attraction often uses recorded vocalizations and decoys to attract new breeders^{95,139,146}. Social attraction has been most successful on species with little natal philopatry and short generation times, such as gulls and terns, or when there are nearby source colonies^{95,147}. A review of seabird restoration projects found that 83% of Laridae social attraction projects with known outcomes have resulted in successful breeding¹⁴⁸.

Many seabirds such as shearwaters, petrels, storm-petrels and albatrosses exhibit strong natal philopatry, which restricts the colonization of new restoration sites, even if social attraction is used^{95,139,140}. For such species, egg or chick translocations are the best way to establish new colonies or increase existing ones. Most translocations involve moving cohorts of chicks from their natal colony to the restoration site before the chicks have imprinted on their natal colony. Translocated chicks then imprint on the restoration site and, upon sexual maturity, return to the restoration site to breed⁹⁵. Of the 851 reported seabird restoration efforts up to 2021, only 6% have included translocation⁹⁵, because of the high cost of translocations, the expertise needed to rear chicks and the lack of identified suitable habitat for colony creation¹⁴⁰. The founding translocated chicks help to socially attract other conspecifics and species. Translocated seabirds might also use new marine foraging grounds based on their new fledge location, which re-establishes or newly establishes a circular seabird economy around the restoration site. Seabird restoration can unite people and communities and increase the cooperation among communities, land managers and conservation practitioners, which also increases the chances of long-term success for the newly founded colony^{139,140,149}.

Most research studies the efficacy of seabird social attraction and translocation itself^{95,143}, rather than the ecosystem-level impacts

of seabird restoration. What little work that has been conducted suggests that seabird translocation results in positive ecosystem impacts, including increased seabird-derived nutrients in island food webs and arthropod diversity¹⁵⁰. Ecosystems also show marked improvements of nutrient cycling, biodiversity and community composition when seabirds passively recover from invasive mammal removal^{29,66,68,136,137,151,152}. However, full restoration of all components of the circular seabird economy could take decades or longer, owing to different recovery times of various island ecosystem constituents¹⁵³. Specifically, invertebrates and faster-growing plants tend to respond fastest to seabird density increases, whereas seabird-derived nutrient cycling and flora–fauna community composition take longer to recover¹⁵³. Because they respond most quickly to restoration attempts, gulls and terns might be conduits of faster ecosystem restoration than species that respond more slowly, such as petrels, shearwaters and albatrosses. More research on which seabird species can most effectively recover the circular seabird economy could help practitioners seeking whole-ecosystem responses.

Restoring seabirds for climate change resilience

Restoring seabird colonies and augmenting existing ones through seabird restoration can help to combat the biodiversity crisis by restoring the circular seabird economy, with all the attendant impacts to islands and oceans. However, seabirds sit at the nexus of both the biodiversity and climate crises, and thus also have the potential to contribute to ecosystem adaptations and resilience against climate change.

The impacts of climate change on seabirds have been extensively reviewed^{9,12,13}, but few direct attempts have been made to determine the impacts of seabirds on climate change resilience (though see refs. 47,104,111). However, the broad-scale and large-scale effects of seabirds on ecosystem functioning suggest likely interconnectedness. Focusing on the major resilience factors and adaptive capacities of ecosystems, and how these are influenced by the circular seabird economy, can guide future research on the interactions between the circular seabird economy and climate resilience⁶⁰. Particularly relevant is the extent to which seabirds improve the adaptive capacity of terrestrial and marine systems to climate-induced shifts (such as temperature, water availability, storm surge, sea level rise and severe weather events), as well as their potential to mitigate climate change via effects on carbon sequestration.

Given that the impacts of both climate change and the circular seabird economy vary across biomes, the relevance of seabirds for climate resilience will certainly vary biogeographically. As described, seabirds have particularly strong effects on nutrient cycling in polar and subpolar regions, where there are limited other external nutrient subsidies^{22,26}. In turn, seabirds have a major role in shaping vegetation and maintaining the functioning of terrestrial and marine communities and ecosystems near their colonies, increasing localized climate resilience when such systems are protected or restored. However, seabirds might also negatively affect climate resilience and mitigation in the short term, especially in ecosystems in which plant communities are in the process of recovery, as seabird burrowing can alter forest regeneration and composition, which accelerates erosion^{68,69,154}.

More is known about the effects of seabirds on climate resilience in the tropics, where low-lying atoll and reef islands are major seabird hotspots^{155,156}. To withstand projected sea level rise, increased storm surges and severe rain events, these islands require deposition of reef-borne materials to maintain their adaptive growth capacity^{148,157,158}, and intact coral reefs that provide a physical barrier

against high-energy oceanic erosive forces^{159,160}. Through their nutrient inputs, seabirds can enhance these resilience factors by contributing to increased coral growth and calcification rates^{103–105}, sediment generation rates from bio-eroding parrotfish⁸⁸ and mangrove ecosystem productivity¹¹⁶. Tropical coral reefs are themselves some of the most vulnerable ecosystems to climate change, as increasing marine heatwaves cause mass die-offs of reef-building corals¹⁶¹. Seabird nutrient inputs contribute directly to coral reef resilience by facilitating faster coral recovery following extreme marine heatwaves¹⁰⁴ and might also indirectly enhance resilience through their effects on other reef organisms (for example, crustose coralline algae and herbivorous fishes)^{88,111}.

There is now increasing recognition that conservation interventions, such as eradicating invasive predators on islands, could also benefit climate resilience via benefits to seabirds, as well as by other direct and indirect effects^{162,163}. Likewise, native vegetation, seabirds and climate resilience are intricately interconnected – with seabirds preferring native vegetation and also benefiting it via nutrient inputs⁴⁴, and native vegetation influencing climate mitigation via carbon stocks and climate adaptive capacity via soil stability¹⁶⁴. Such feedback between climate change impacts on seabirds, and seabird impacts on climate

change resilience, suggests that holistic ecosystem restoration could be an avenue to provide mutualistic benefits.

Summary and future directions

In Hawai'i, early naturalist observations, palaeontological, genetic and isotopic evidence, demonstrate 'ua'u or Hawaiian petrel were once widespread across the archipelago, and foraged at a higher trophic level in the pelagic zone^{165–167}. On atolls, where comparable historical evidence is often intractable, seabird diversity and abundance thrives where anthropogenic threats are minimal, indicating seabird impacts globally were previously more widespread^{88,168}. Thus, historically, seabirds were vastly more abundant, and circular seabird economies connecting oceans and islands were larger in magnitude and complexity than they are today. Despite global seabird declines, this Review presents the large-scale, cross-ecosystem impacts seabirds have both in oceans and on their associated islands. Seabirds have top predator roles in oceans, transfer nutrients to islands, resulting in unique biogeochemistry and ecology, both on islands and in surrounding marine communities. Even though seabirds continue to face numerous threats, restoration efforts have been largely successful; further restoration can ensure that the circular seabird economy continues

Box 2 | Emerging approaches to explore the circular seabird economy

The impacts of the circular seabird economy are far-reaching. Identifying and measuring useful indicators of these impacts are subject to time and logistical constraints, availability of expertise and technological limitations. However, the introduction and novel application of methods, including energetic approaches, eco-acoustics, remote sensing and omics techniques, is continually improving data collection capabilities and growing scientists' mechanistic, spatial and temporal understanding of the circular seabird economy.

Energy requirements, which underpin all biological and ecological interactions, have been largely neglected in conservation biology despite being a well-established field. The application of ecological energetics allows exploration of the influence of resource availability and use on organismal metabolisms and species interactions in various ecosystems and compares disparate organisms using a common currency¹⁹⁹. Indeed, energetic approaches have been used to evaluate the influence of land-use changes on species diversity and dynamics²⁰⁰ and to project the influence of tropical island restoration for seabirds and coral reefs¹⁹⁴. Given that seabird energetics are inextricably linked to nutrient flows between trophic levels and across ecosystems, the predictive capacity of ecological energetic modelling can contribute useful indicators of a healthy seabird economy and support informed island management practices.

Remote-sensing methodologies provide additional opportunities to measure indicators over greater spatial and temporal scales, with higher sampling frequency and resolution. Non-invasive tools such as camera traps, drones and radar are already used to record seabird biomass and abundances^{201,202}. Satellite images and Earth observation data that use hyperspectral or multispectral sensors can help to determine both direct and indirect seabird impacts on islands and nearshore habitats (reviewed elsewhere²⁰³). For example, seabird

guano has a distinct spectral signature that can be used to locate seabird colonies²⁰⁴, and spectral, infrared, chlorophyll *a* or turbidity sensors can identify nutrients in marine habitats^{205,206}, vegetation canopies^{207–210} and the atmosphere²¹¹. Coupled with machine-learning or artificial intelligence approaches²¹², automated analyses of vast amounts of imagery and remote-sensing data can be performed while reducing data-accessibility barriers. With increased availability and lowered cost of tools such as recording devices and analytical software^{213,214}, eco-acoustic monitoring has proven effective in non-invasively recording seabird biodiversity^{215,216}, colony density¹⁶⁵ and population recovery following seabird restoration efforts¹⁶⁶. Eco-acoustics can also support measurements of cross-ecosystem impacts, as they are known to provide proxies for marine biodiversity¹⁶⁷ and ecosystem recovery²¹⁷.

The use of omics tools that rely on DNA or RNA in environmental or biological samples enables the identification of species presence or absence, relative abundance and activity of entire land-sea assemblages within areas influenced by seabirds. For example, environmental DNA can be used to estimate seabird biodiversity²¹⁸, understand seabird diet²¹⁹ or analyse ancient sedimentary DNA to recover historic seabird baselines²²⁰. Researchers can also use environmental DNA to detect invasive predators that might threaten seabird populations^{221,222}, and explore shifts or variability in entire marine communities^{223,224}. Similarly, 16S rRNA metabarcoding can reveal the impact of seabird nutrient subsidies on microbial community dynamics in island soils²²⁵ and nearshore marine habitats¹¹², offering early biological indicators of broader ecological change²²⁶. Additional (meta)genomics and (meta)transcriptomics tools could further elucidate taxa and genes associated with seabird-derived nutrient uptake to help to identify indicator species that might support a more mechanistic approach to ecosystem monitoring²²⁷.

to thrive in the face of large-scale ecosystem change. In an ideal world, the seabird restoration techniques discussed in this Review could be the conduit for repairing lost linkages between oceans and islands globally. Indeed, if seabird restoration reaches its fullest potential, all sparsely or uninhabited islands would be rich in seabirds and the benefits they provide.

To restore the circular seabird economy at scale, there are important knowledge gaps that should be addressed regarding the recovery of nutrient cycling and ecosystem function during seabird recovery. These gaps can be filled using traditional and emerging approaches (Box 2). Seabird-derived nutrients, such as nitrogen, have measurable declines with time since seabird colony abandonment¹⁶⁹. Although these nutrients often increase in the decades following invasive species removal^{136,170}, a historical baseline of isotopic enrichment or nutrient availability to evaluate recovery is rarely available, though the use of nearby, relatively undisturbed islands can be useful recovery targets^{136,150,171}. Additionally, it remains unclear when islands no longer function as seabird islands – that is, when terrestrial island systems are no longer driven by seabird-derived subsidies. Seabird-derived nutrients can be highly localized in terrestrial systems, declining with distance from colonies¹⁷², and with island size¹⁵¹. Further research on potential island size thresholds beyond which seabirds do not dominate nutrient cycling, or ecosystem circumstances that may prevent reinstatement of the circular seabird economy, is necessary. Ideally, researching seabird nutrient dynamics and island size thresholds would be done at scale throughout various biomes, to improve understanding of how patterns change across different ecosystems. The technology to track seabird impacts at scale is emerging (Box 2), and as it expands, so too will researchers' ability to tackle knowledge gaps globally.

Cross-system and marine effects began to be systematically studied around a decade ago^{36,173–175}. Accordingly, knowledge about seabird nutrient effects is patchy, and several critical habitats such as mangroves, reefs, oyster beds and seagrass beds are understudied even when seabird benefits have been established. The partial or total absence of replication for some of these systems further limits researchers' ability to reliably quantify and predict seabird effects and how these interact with local conditions and stressors. Likewise, cross-system studies integrating both terrestrial and marine habitats remain scarce, representing 5% of all seabird effect publications²⁶, offering a poor understanding of the timescales over which terrestrial and marine outcomes return following seabird restoration¹⁵³. This scarcity is in spite of extensive evidence of strong ecological connectivity between islands, coastal areas and the surrounding nearshore marine habitats, especially for atolls⁶⁰.

Beyond habitats, this patchiness extends to marine organisms, with very little understanding of how seabird nutrients affect marine consumers³⁶. Likewise, there is a paucity of work on how seasonal and temporal variability affects seabird nutrient delivery (but see ref. 172). Importantly, the lack of systematic and replicated documentation of seabird effects in marine environments, both with respect to nearshore impacts and the link between nearshore impacts and the pelagic ecosystems in which most seabirds forage (for example, ref. 93), directly undermines the potential of these studies for building knowledge, informing decision-makers and guiding conservation practices that target marine environments¹⁶⁸. Although this Review has mostly focused on regional scale impacts, it is important to note that many seabirds move within large global marine flyways¹⁷⁶, suggesting nutrient and energy flows with land–sea impacts at a global scale similar to

other mobile marine vertebrates (for example, ref. 177) that have rarely been studied for seabirds. Future research must address these spatial and temporal gaps by fostering a more integrated marine-terrestrial, and planetary, perspective.

Gaps in knowledge about seabirds and their interactions have complex implications, which are probably underestimated in dominant forms of science. Seabirds provide both direct and indirect benefits to people and have a central role in many Indigenous cultures and local communities around the world, particularly given that coastal and island communities rely on both subsistence and market economies for their livelihoods¹⁴⁹. For example, 80% of Maldivian tuna fishers navigate and locate fishing grounds by following seabirds¹⁷⁸. Seabirds provide provisioning, supportive and cultural services with sizeable historical connections to human sustenance and local economies in which communities mined seabird guano as a fertilizer for agriculture and harvested seabirds and their eggs for food¹⁷⁹. The benefits of the circular seabird economy to people largely focus on the indirect impact that their supportive services (nutrient cycling, seed dispersal and ecosystem engineering) have to the ecosystems that people rely on – such as increased fish biomass⁸⁸, improved coastal ecosystem resilience¹⁰⁴ and carbon sequestration¹⁶³.

Moreover, dominant forms of science have eschewed Indigenous and traditional ecological knowledge, potentially placing conservation actions such as restricting seabird harvesting at odds with Indigenous resource management practices¹⁴⁹. Future research into the benefits of the circular seabird economy should seek a more holistic, people-centred approach that integrates Indigenous ways of knowing, perspectives and cultural benefits.

Published online: 27 October 2025

References

- Brooke, M. The food consumption of the world's seabirds. *Proc. R. Soc. Lond. B Biol. Sci.* **271**, S246–S248 (2004).
- Karpouzli, V. S., Watson, R. & Pauly, D. Modelling and mapping resource overlap between seabirds and fisheries on a global scale: a preliminary assessment. *Mar. Ecol. Prog. Ser.* **343**, 87–99 (2007).
- Grémillet, D. et al. Persisting worldwide seabird-fishery competition despite seabird community decline. *Curr. Biol.* **28**, 4009–4013.e2 (2018).
- OECD. *OECD Review of Fisheries 2025* (OECD Publishing, 2025).
- Schreiber, E. A. & Burger, J. *Biology of Marine Birds* (CRC Press, 2001).
- Jovani, R. et al. Colony size and foraging range in seabirds. *Oikos* **125**, 968–974 (2016).
- Mulder, C. P. H. et al. (eds) *Seabird Islands: Ecology, Invasion, and Restoration* (Oxford Univ. Press, 2011).
- Jones, H. P., Borrelle, S. B. & Rankin, L. L. Land–sea linkages depend on macroalgal species, predator invasion history in a New Zealand archipelago. *Restor. Ecol.* **31**, e13798 (2023).
- Dias, M. P. et al. Threats to seabirds: a global assessment. *Biol. Conserv.* **237**, 525–537 (2019).
- Fernández-Palacios, J. M. et al. Scientists' warning — the outstanding biodiversity of islands is in peril. *Glob. Ecol. Conserv.* **31**, e01847 (2021).
- Borrelle, S., Jones, H., Richard, Y. & Salguero-Gomez, R. Estimating the impact of marine threats to seabird recovery after predator eradication. *Mar. Ornithol.* **51**, 225–236 (2023).
- Sydeman, W. J., Thompson, S. A. & Kitaysky, A. Seabirds and climate change: roadmap for the future. *Mar. Ecol. Prog. Ser.* **454**, 107–117 (2012).
- Orgeret, F. et al. Climate change impacts on seabirds and marine mammals: the importance of study duration, thermal tolerance and generation time. *Ecol. Lett.* **25**, 218–239 (2022).
- Croxall, J. P. et al. Seabird conservation status, threats and priority actions: a global assessment. *Bird. Conserv. Int.* **22**, 1–34 (2012).
- Jodice, P. G. R. & Suryan, R. M. in *Landscape-Scale Conservation Planning* (eds Trombulak, S. C. & Baldwin, R. F.) 139–165 (Springer Netherlands, 2010).
- Dunn, R. E., White, C. R. & Green, J. A. A model to estimate seabird field metabolic rates. *Biol. Lett.* **14**, 20180190 (2018).
- Allen, P., Peters, A., Stiefel, Z. & Clarke, R. H. Seabird moult timing and duration: implications for at-sea threat exposure worldwide. *Glob. Ecol. Conserv.* **59**, e03558 (2025).
- Smith, J. L., Mulder, C. P. H. & Ellis, J. C. in *Seabird Islands: Ecology, Invasion, and Restoration* (eds Mulder, C. P. H. et al.) 27–55 (Oxford Univ. Press, 2011).

19. De La Peña-Lastra, S., Pérez-Alberti, A., Ferreira, T. O., Huerta-Díaz, M. Á & Otero, X. L. Global deposition of potentially toxic metals via faecal material in seabird colonies. *Sci. Rep.* **12**, 22392 (2022).
20. Phillips, R. A., Lewis, S., González-Solís, J. & Daunt, F. Causes and consequences of individual variability and specialization in foraging and migration strategies of seabirds. *Mar. Ecol. Prog. Ser.* **578**, 117–150 (2017).
21. Bird, M. I., Tait, E., Wurster, C. M. & Furness, R. W. Stable carbon and nitrogen isotope analysis of avian uric acid. *Rapid Commun. Mass Spectrom.* **22**, 3393–3400 (2008).
22. Otero, X. L., De La Peña-Lastra, S., Pérez-Alberti, A., Ferreira, T. O. & Huerta-Díaz, M. A. Seabird colonies as important global drivers in the nitrogen and phosphorus cycles. *Nat. Commun.* **9**, 246 (2018).
23. Riddick, S. N. et al. Global assessment of the effect of climate change on ammonia emissions from seabirds. *Atmos. Environ.* **184**, 212–223 (2018).
24. Schmidt, S. et al. Atmospheric concentrations of ammonia and nitrogen dioxide at a tropical coral cay with high seabird density. *J. Environ. Monit.* **12**, 460–465 (2010).
25. Wing, S. R. et al. Seabirds and marine mammals redistribute bioavailable iron in the Southern Ocean. *Mar. Ecol. Prog. Ser.* **510**, 1–13 (2014).
26. Grant, M. L., Bond, A. L. & Lavers, J. L. The influence of seabirds on their breeding, roosting and nesting grounds: a systematic review and meta-analysis. *J. Anim. Ecol.* **91**, 1266–1289 (2022).
27. Roman, L. et al. Nutrients and seabird biogeography: feather elements differ among oceanic basins in the Southern Hemisphere, reflecting bird size, foraging range and nutrient availability in seawater. *Glob. Ecol. Biogeogr.* **32**, 495–510 (2023).
28. Grant, M. L., Bond, A. L., Reichman, S. M. & Lavers, J. L. Seabird transported contaminants are dispersed in island ecosystems. *Chemosphere* **361**, 142483 (2024).
29. Benkwitt, C. E., Gunn, R. L., Le Corre, M., Carr, P. & Graham, N. A. Rat eradication restores nutrient subsidies from seabirds across terrestrial and marine ecosystems. *Curr. Biol.* **31**, 2704–2711 (2021).
30. Healing, S., Benkwitt, C. E., Dunn, R. E. & Graham, N. A. Seabird-vectored pelagic nutrients integrated into temperate intertidal rocky shores. *Front. Mar. Sci.* **11**, 1343966 (2024).
31. Rodgers, K. A. The cycle of phosphorus in an atoll ecosystem. *Appl. Geochem.* **9**, 311–322 (1994).
32. McMahon, A. & Santos, I. R. Nitrogen enrichment and speciation in a coral reef lagoon driven by groundwater inputs of bird guano. *J. Geophys. Res. Ocean.* **122**, 7218–7236 (2017).
33. Riddick, S. N. et al. The global distribution of ammonia emissions from seabird colonies. *Atmos. Environ.* **55**, 319–327 (2012).
34. Wentworth, G. R. et al. Ammonia in the summertime Arctic marine boundary layer: sources, sinks, and implications. *Atmos. Chem. Phys.* **16**, 1937–1953 (2016).
35. Zwolicki, A., Zmudczyńska-Skarbek, K. M., Iliszko, L. & Stempniewicz, L. Guano deposition and nutrient enrichment in the vicinity of planktivorous and piscivorous seabird colonies in Spitsbergen. *Polar Biol.* **36**, 363–372 (2013).
36. De La Peña-Lastra, S. Seabird droppings: effects on a global and local level. *Sci. Total Environ.* **754**, 142148 (2021).
37. Benkwitt, C. E., Carr, P., Wilson, S. K. & Graham, N. A. Seabird diversity and biomass enhance cross-ecosystem nutrient subsidies. *Proc. R. Soc. B* **289**, 20220195 (2022).
38. Appoo, J. et al. Seabird presence and seasonality influence nutrient dynamics of atoll habitats. *Biotropica* **56**, e13354 (2024).
39. Votier, S. C. et al. Geolocation and immersion loggers reveal year-round residency and facilitate nutrient deposition rate estimation of adult red-footed boobies in the Chagos Archipelago, tropical Indian Ocean. *J. Avian Biol.* **2024**, e03185 (2024).
40. Rankin, L. L. & Jones, H. P. Nearshore ecosystems on seabird islands are potentially influenced by invasive predator eradications and environmental conditions: a case study at the Mercury Islands, New Zealand. *Mar. Ecol. Prog. Ser.* **661**, 83–96 (2021).
41. Sandin, S. A. et al. Harnessing island–ocean connections to maximize marine benefits of island conservation. *Proc. Natl Acad. Sci. USA* **119**, e2122354119 (2022).
42. Stapp, P., Polls, G. A. & Pinero, F. S. Stable isotopes reveal strong marine and El Niño effects on island food webs. *Nature* **401**, 467–469 (1999).
43. Roos, R. E. et al. Marine-derived nutrients shape the functional composition of high arctic plant communities. *Funct. Ecol.* **9**, 1606–1621 (2025).
44. Young, H. S., McCauley, D. J., Dunbar, R. B. & Dirzo, R. Plants cause ecosystem nutrient depletion via the interruption of bird-derived spatial subsidies. *Proc. Natl Acad. Sci. USA* **107**, 2072–2077 (2010).
45. Appoo, J., Graham, N. A. J., Jones, C. W., Jaquemet, S. & Bunbury, N. Seabird nutrient subsidy alters size and resource use of functionally important mangrove macroinvertebrates. *Ecosphere* **15**, e70121 (2024).
46. Wu, L. et al. Nitrogen cycling in the soil–plant system along a series of coral islands affected by seabirds in the South China Sea. *Sci. Total Environ.* **627**, 166–175 (2018).
47. Wardle, D. A., Bellingham, P. J., Fukami, T. & Mulder, C. P. H. Promotion of ecosystem carbon sequestration by invasive predators. *Biol. Lett.* **3**, 479–482 (2007).
48. Orwin, K. H. et al. Burrowing seabird effects on invertebrate communities in soil and litter are dominated by ecosystem engineering rather than nutrient addition. *Oecologia* **180**, 217–230 (2016).
49. Ellis, J. C. Marine birds on land: a review of plant biomass, species richness, and community composition in seabird colonies. *Plant Ecol.* **181**, 227–241 (2005).
50. Sanchez-Pinero, F. & Polis, G. A. Bottom-up dynamics of allochthonous input: direct and indirect effects of seabirds on islands. *Ecology* **81**, 3117–3132 (2000).
51. Leblans, N. I. W. et al. Effects of seabird nitrogen input on biomass and carbon accumulation after 50 years of primary succession on a young volcanic island, Surtsey. *Biogeosciences* **11**, 6237–6250 (2014).
52. Duda, M. P. et al. Long-term changes in terrestrial vegetation linked to shifts in a colonial seabird population. *Ecosystems* **23**, 1643–1656 (2020).
53. Croll, D. A., Maron, J. L., Estes, J. A., Danner, E. M. & Byrd, G. V. Introduced predators transform subarctic islands from grassland to tundra. *Science* **307**, 1959–1961 (2005).
54. Gillham, M. E. Vegetation of sea and shore-bird colonies on Aldabra atoll. *Atoll Res. Bull.* **200**, 1–19 (1977).
55. Ellis, J. C., Fariña, J. M. & Witman, J. D. Nutrient transfer from sea to land: the case of gulls and cormorants in the Gulf of Maine. *J. Animal Ecol.* **75**, 565–574 (2006).
56. Maron, J. L. et al. An introduced predator alters Aleutian Island plant communities by thwarting nutrient subsidies. *Ecol. Monogr.* **76**, 3–24 (2006).
57. Bancroft, W. J., Garkaklis, M. J. & Roberts, J. D. Burrow building in seabird colonies: a soil-forming process in island ecosystems. *Pedobiologia* **49**, 149–165 (2005).
58. Wait, D. A., Aubrey, D. P. & Anderson, W. B. Seabird guano influences on desert islands: soil chemistry and herbaceous species richness and productivity. *J. Arid Environ.* **60**, 681–695 (2005).
59. Young, H. S., McCauley, D. J. & Dirzo, R. Differential responses to guano fertilization among tropical tree species with varying functional traits. *Am. J. Bot.* **98**, 207–214 (2011).
60. Steibl, S., Bunbury, N., Young, H. S. & Russell, J. C. A renaissance of atoll ecology. *Annu. Rev. Ecol. Syst.* **55**, 301–322 (2024).
61. Burger, A. E. Dispersal and germination of seeds of *Pisonia grandis*, an Indo-Pacific tropical tree associated with insular seabird colonies. *J. Trop. Ecol.* **21**, 263–271 (2005).
62. Calvino-Cancela, M. P. & Anderson, W. B. Seabird guano influences on desert islands: soil chemistry and herbaceous species richness and productivity. *Plant Ecol.* **212**, 1149–1157 (2011).
63. Aoyama, Y., Kawakami, K. & Chiba, S. Seabirds as adhesive seed dispersers of alien and native plants in the oceanic Ogasawara Islands, Japan. *Biodivers. Conserv.* **21**, 2787–2801 (2012).
64. Mulder, C. P. H., Wardle, D. A., Durrett, M. S. & Bellingham, P. J. Leaf damage by herbivores and pathogens on New Zealand islands that differ in seabird densities. *N. Z. J. Ecol.* **39**, 221–230 (2015).
65. Hawke, D. J. & Vallance, J. R. Microbial carbon concentration in samples of seabird and non-seabird forest soil: implications for leaf litter cycling. *Pedobiologia* **58**, 33–39 (2015).
66. Fukami, T. et al. Above-and below-ground impacts of introduced predators in seabird-dominated island ecosystems. *Ecol. Lett.* **9**, 1299–1307 (2006).
67. Hayashi, K. et al. Seabird-affected taluses are denitrification hotspots and potential N₂O emitters in the high arctic. *Sci. Rep.* **8**, 17261 (2018).
68. Grant-Hoffman, M. N., Mulder, C. P. & Bellingham, P. J. Invasive rats alter woody seedling composition on seabird-dominated islands in New Zealand. *Oecologia* **163**, 449–460 (2010).
69. Rodway, M. S., Wilson, L. K., Lemon, M. J. & Millikin, R. L. The ups and downs of ecosystem engineering by burrow-nesting seabirds on Triangle Island, British Columbia. *Mar. Ornithol.* **45**, 47–55 (2017).
70. Towns, D. R., Daugherty, C. H. & Cree, A. Raising the prospects for a forgotten fauna: a review of 10 years of conservation effort for New Zealand reptiles. *Biol. Conserv.* **99**, 3–16 (2001).
71. Markwell, T. J. & Daugherty, C. H. Invertebrate and lizard abundance is greater on seabird-inhabited islands than on seabird-free islands in the Marlborough Sounds, New Zealand. *Ecoscience* **9**, 293–299 (2002).
72. Bellingham, P. J. et al. New Zealand island restoration: seabirds, predators, and the importance of history. *N. Z. J. Ecol.* **34**, 115 (2010).
73. Polis, G. A. & Hurd, S. D. Linking marine and terrestrial food webs: allochthonous input from the ocean supports high secondary productivity on small islands and coastal land communities. *Am. Nat.* **147**, 396–423 (1996).
74. Ellis, J. C. et al. in *Seabird Islands: Ecology, Invasion, and Restoration* (eds Mulder, C. P. H. et al.) 177–211 (Oxford Univ. Press, 2011).
75. Kolb, G. S., Young, H. S. & Anderson, W. B. in *Seabird Islands: Ecology, Invasion, and Restoration* (eds Mulder, C. P. H. et al.) 212–241 (Oxford Univ. Press, 2011).
76. Halpin, L. R. et al. Arthropod predation of vertebrates structures trophic dynamics in island ecosystems. *Am. Nat.* **198**, 540–550 (2021).
77. Lane, A. M. & Shine, R. When seasnake meets seabird: ecosystem engineering, facilitation and competition. *Austral Ecol.* **36**, 544–549 (2011).
78. Corkery, I., Bell, B. & Nelson, N. Is the breeding behaviour of nesting seabirds influenced by the presence of a predatory reptile — the tuatara? *J. R. Soc. N. Z.* **45**, 21–30 (2015).
79. Fischer, J. H., McCauley, C. F., Armstrong, D. P., Debski, I. & Wittmer, H. U. Contrasting responses of lizard occurrences to burrowing by a critically endangered seabird. *Community Ecol.* **20**, 64–74 (2019).
80. Towns, D. *Ahuahu: An Island Conservation Journey in Aotearoa New Zealand* (Canterbury Univ. Press, 2023).
81. Whitworth, D. L., Carter, H. R. & Gress, F. Recovery of a threatened seabird after eradication of an introduced predator: eight years of progress for Scripps's murrelet at Anacapa Island, California. *Biol. Conserv.* **162**, 52–59 (2013).
82. Watson, J., Leitch, A. F. & Broad, R. A. The diet of the sea eagle *Haliaeetus albicilla* and golden eagle *Aquila chrysaetos* in western Scotland. *Ibis* **134**, 27–31 (1992).
83. Brattstrom, B. H. Food webs and feeding habits on the Revillagigedo Islands, Mexico. *Pac. Sci.* **69**, 181–195 (2015).
84. Mills, K. L. Seabirds as part of migratory owl diet on Southeast Farallon Island, California. *Mar. Ornithol.* **44**, 121–126 (2016).
85. Whitehead, E. A. Little shearwaters (*Puffinus assimilis haurakiensis*) as prey for morepork (*Ninox novaeseelandiae*). *Notornis* **68**, 170–172 (2021).
86. Bosman, A. L. & Hockey, P. A. R. Seabird guano as a determinant of rocky intertidal community structure. *Mar. Ecol. Prog. Ser.* **32**, 247–257 (1986).

87. Kurle, C. M., Croll, D. A. & Tershy, B. R. Introduced rats indirectly change marine rocky intertidal communities from algae-to invertebrate-dominated. *Proc. Natl Acad. Sci. USA* **105**, 3800–3804 (2008).
88. Graham, N. A. J. et al. Seabirds enhance coral reef productivity and functioning in the absence of invasive rats. *Nature* **559**, 250–253 (2018).
89. Benkwitt, C. E., Wilson, S. K. & Graham, N. A. Biodiversity increases ecosystem functions despite multiple stressors on coral reefs. *Nat. Ecol. Evol.* **4**, 919–926 (2020).
90. Moss, B. Marine reptiles, birds and mammals and nutrient transfers among the seas and the land: an appraisal of current knowledge. *J. Exp. Mar. Biol. Ecol.* **492**, 63–80 (2017).
91. Dunn, R. E. et al. Active and passive pathways of nutrient transfer in coral reef ecosystems. *Coral Reefs* <https://doi.org/10.1007/s00338-025-02676-z> (2025).
92. Cumming, G. S., James, N. L., Chua, C. M. & Huertas, V. A framework and review of evidence of the importance of coral reefs for marine birds in tropical ecosystems. *Ecol. Evol.* **14**, e70165 (2024).
93. Weber, S. B. et al. Direct evidence of a prey depletion ‘halo’ surrounding a pelagic predator colony. *Proc. Natl Acad. Sci. USA* **118**, e2101325118 (2021).
94. Unsworth, R. K. & Butterworth, E. G. Seagrass meadows provide a significant resource in support of avifauna. *Diversity* **13**, 363 (2021).
95. Spatz, D. R. et al. Tracking the global application of conservation translocation and social attraction to reverse seabird declines. *Proc. Natl Acad. Sci. USA* **120**, e2214574120 (2023).
96. Shatova, O., Wing, S. R., Gault-Ringold, M., Wing, L. & Hoffmann, L. J. Seabird guano enhances phytoplankton production in the Southern Ocean. *J. Exp. Mar. Biol. Ecol.* **483**, 74–87 (2016).
97. Alba-González, P., Álvarez-Salgado, X. A., Cobelo-García, A., Kaal, J. & Teira, E. Faeces of marine birds and mammals as substrates for microbial plankton communities. *Mar. Environ. Res.* **174**, 105560 (2022).
98. Browning, T. J. et al. The role of seabird guano in maintaining North Atlantic summertime productivity. *Sci. Total Environ.* **897**, 165309 (2023).
99. Bosman, A. L., Du Toit, J. T., Hockey, P. A. R. & Branch, G. M. A field experiment demonstrating the influence of seabird guano on intertidal primary production. *Estuar. Coast. Shelf Sci.* **23**, 283–294 (1986).
100. Powell, G. V., Fourqurean, J. W., Kenworthy, W. J. & Zieman, J. C. Bird colonies cause seagrass enrichment in a subtropical estuary: observational and experimental evidence. *Estuar. Coast. Shelf Sci.* **32**, 567–579 (1991).
101. Young, H. S., Hurrey, L. & Kolb, G. S. in *Seabird Islands: Ecology, Invasion, and Restoration* (eds Mulder, C. P. H. et al.) 242–260 (Oxford Univ. Press, 2011).
102. Kazama, K. Bottom-up effects on coastal marine ecosystems due to nitrogen input from seabird feces. *Ornithol. Sci.* **18**, 117–126 (2019).
103. Savage, C. Seabird nutrients are assimilated by corals and enhance coral growth rates. *Sci. Rep.* **9**, 4284 (2019).
104. Benkwitt, C. E. et al. Seabirds boost coral reef resilience. *Sci. Adv.* **9**, ead0390 (2023).
105. Lange, I. D. & Benkwitt, C. E. Seabird nutrients increase coral calcification rates and boost reef carbonate production. *Sci. Rep.* **14**, 24937 (2024).
106. Jeannot, L.-L., Lozano-Peña, J., Zora, A., Brandl, S. & Graham, N. A. J. Seabird-derived nutrients influence feeding pathways and body size in cryptobenthic reef fishes. *Proc. R. Soc. B* **292**, 20250539 (2025).
107. Wootton, J. T. Direct and indirect effects of nutrients on intertidal community structure: variable consequences of seabird guano. *J. Exp. Mar. Biol. Ecol.* **151**, 139–153 (1991).
108. Methratta, E. T. Top-down and bottom-up factors in tidepool communities. *J. Exp. Mar. Biol. Ecol.* **299**, 77–96 (2004).
109. Littler, M. M., Littler, D. S. & Tillyanov, E. A. Comparisons of N- and P-limited productivity between high granitic islands versus low carbonate atolls in the Seychelles Archipelago: a test of the relative-dominance paradigm. *Coral Reefs* **10**, 199–209 (1991).
110. Lapointe, B. E., Littler, M. M. & Littler, D. S. Modification of benthic community structure by natural eutrophication: the Belize barrier reef. In *Proc. Seventh International Coral Reefs Symposium, Guam, 1992*, vol. 1, 323–334 (Univ. Guam Marine Laboratory, 1993).
111. Benkwitt, C. E., Wilson, S. K. & Graham, N. A. J. Seabird nutrient subsidies alter patterns of algal abundance and fish biomass on coral reefs following a bleaching event. *Glob. Change Biol.* **25**, 2619–2632 (2019).
112. Benkwitt, C. E. et al. Re-connecting ecosystems: integrating coral reefs into monitoring of island restoration. *Ecol. Indic.* **170**, 113042 (2025).
113. Fourqurean, J. W., Powell, G. V., Kenworthy, W. J. & Zieman, J. C. The effects of long-term manipulation of nutrient supply on competition between the seagrasses *Thalassia testudinum* and *Halodule wrightii* in Florida Bay. *Oikos* **72**, 349–358 (1995).
114. Shatova, O. A., Wing, S. R., Hoffmann, L. J., Wing, L. C. & Gault-Ringold, M. Phytoplankton community structure is influenced by seabird guano enrichment in the Southern Ocean. *Estuar. Coast. Shelf Sci.* **191**, 125–135 (2017).
115. Hentati-Sundberg, J. et al. Fueling of a marine-terrestrial ecosystem by a major seabird colony. *Sci. Rep.* **10**, 15455 (2020).
116. Appoo, J., Bunbury, N., Jaquemet, S. & Graham, N. A. Seabird nutrient subsidies enrich mangrove ecosystems and are exported to nearby coastal habitats. *iScience* **27**, 109404 (2024).
117. Andrades, R. et al. Seabird guano reshapes intertidal reef food web in an isolated oceanic islet. *Coral Reefs* **43**, 347–355 (2024).
118. Onuf, C. P., Teal, J. M. & Valiela, I. Interactions of nutrients, plant growth and herbivory in a mangrove ecosystem. *Ecology* **58**, 514–526 (1977).
119. Vizzini, S., Signa, G. & Mazzola, A. Guano-derived nutrient subsidies drive food web structure in coastal ponds. *PLoS ONE* **11**, e0151018 (2016).
120. Kolb, G. S., Ekholm, J. & Hambäck, P. A. Effects of seabird nesting colonies on algae and aquatic invertebrates in coastal waters. *Mar. Ecol. Prog. Ser.* **417**, 287–300 (2010).
121. Gagnon, K., Rothäusler, E., Syrjänen, A., Yli-Renko, M. & Jormalainen, V. Seabird guano fertilizes Baltic Sea littoral food webs. *PLoS ONE* **8**, e61284 (2013).
122. Thibault, M. et al. Seabird-derived nutrients supply modulates the trophic strategies of mixotrophic corals. *Front. Mar. Sci.* **8**, 790408 (2022).
123. Gunn, R. L. et al. Terrestrial invasive species alter marine vertebrate behaviour. *Nat. Ecol. Evol.* **7**, 82–91 (2023).
124. Van Der Plas, F. Biodiversity and ecosystem functioning in naturally assembled communities. *Biol. Rev.* **94**, 1220–1245 (2019).
125. Justel-Diez, M. et al. Inputs of seabird guano alter microbial growth, community composition and the phytoplankton–bacterial interactions in a coastal system. *Environ. Microbiol.* **25**, 1155–1173 (2023).
126. Cardinale, B. J., Ives, A. R. & Inchausti, P. Effects of species diversity on the primary productivity of ecosystems: extending our spatial and temporal scales of inference. *Oikos* **104**, 437–450 (2004).
127. Brose, U. & Hillebrand, H. Biodiversity and ecosystem functioning in dynamic landscapes. *Philos. Trans. R. Soc. B Biol. Sci.* **371**, 20150267 (2016).
128. Thompson, P. L., Isbell, F., Loreau, M., O’Connor, M. I. & Gonzalez, A. The strength of the biodiversity–ecosystem function relationship depends on spatial scale. *Proc. R. Soc. B Biol. Sci.* **285**, 20180038 (2018).
129. Young, L. & VanderWerf, E. (eds) *Conservation of Marine Birds* (Academic Press, 2023).
130. González Ortiz, A. A. et al. Fisheries disrupt marine nutrient cycles through biomass extraction. *Commun. Earth Environ.* **6**, 277 (2025).
131. Rodríguez, A. et al. Artificial lights and seabirds: is light pollution a threat for the threatened Balearic petrels? *J. Ornithol.* **156**, 893–902 (2015).
132. Grant, M. L., Lavers, J. L., Hutton, I. & Bond, A. L. Seabird breeding islands as sinks for marine plastic debris. *Environ. Pollut.* **276**, 116734 (2021).
133. De Jersey, A. M. et al. Seabirds in crisis: plastic ingestion induces proteomic signatures of multiorgan failure and neurodegeneration. *Sci. Adv.* **11**, eads0834 (2025).
134. Clark, B. L. et al. Global assessment of marine plastic exposure risk for oceanic birds. *Nat. Commun.* **14**, 3665 (2023).
135. Garthe, S. et al. Large-scale effects of offshore wind farms on seabirds of high conservation concern. *Sci. Rep.* **13**, 4779 (2023).
136. Jones, H. P. Seabird islands take mere decades to recover following rat eradication. *Ecol. Appl.* **20**, 2075–2080 (2010).
137. Jones, H. P. et al. in *Seabird Islands: Ecology, Invasion, and Restoration* (eds Mulder, C. P. H. et al.) 317–357 (Oxford Univ. Press, 2011).
138. West, J. A. & Nilsson, R. J. Habitat use and burrow densities of burrow-nesting seabirds on South East Island, Chatham Islands, New Zealand. *Notornis* **41**, 27–37 (1994).
139. Jones, H. P. & Kress, S. W. A review of the world’s active seabird restoration projects. *J. Wildl. Manag.* **76**, 2–9 (2012).
140. VanderWerf, E. A. et al. in *Conservation of Marine Birds* (eds Young, L. & VanderWerf, E.) 545–577 (Academic Press, 2023).
141. Major, H. L. & Jones, I. L. An experimental study of the use of social information by prospecting nocturnal burrow-nesting seabirds. *Condor* **113**, 572–580 (2011).
142. Buxton, R. T. & Jones, I. L. An experimental study of social attraction in two species of storm-petrel by acoustic and olfactory cues. *Condor* **114**, 733–743 (2012).
143. Friesen, M. R., Beggs, J. R. & Gaskett, A. C. Sensory-based conservation of seabirds: a review of management strategies and animal behaviours that facilitate success. *Biol. Rev.* **92**, 1769–1784 (2017).
144. Wails, C. N. & Major, H. L. Fitting in with the crowd: the role of prospecting in seabird behavioural trends. *Can. J. Zool.* **95**, 247–253 (2017).
145. Momberg, M. et al. Factors determining nest-site selection of surface-nesting seabirds: a case study on the world’s largest pelagic bird, the wandering albatross (*Diomedea exulans*). *Ibis* **165**, 190–203 (2023).
146. Fischer, J. H., Taylor, G., Debski, I. & Wittmer, H. Acoustic attraction system draws in competing seabird species. *Notornis* **67**, 568–572 (2020).
147. Buxton, R. T., Jones, C., Moller, H. & Towns, D. R. Drivers of seabird population recovery on New Zealand islands after predator eradication. *Conserv. Biol.* **28**, 333–344 (2014).
148. Masselink, G., McCall, R., Beetham, E., Kench, P. & Storlazzi, C. Role of future reef growth on morphological response of coral reef islands to sea-level rise. *J. Geophys. Res. Earth Surf.* **126**, e2020JF005749 (2021).
149. Winter, K. B., Young, R. C. & Lyver, P. O. in *Conservation of Marine Birds* (eds Young, L. & VanderWerf, E.) 321–344 (Academic Press, 2023).
150. Jones, H. P. Prognosis for ecosystem recovery following rodent eradication and seabird restoration in an island archipelago. *Ecol. Appl.* **20**, 1204–1216 (2010).
151. Pascoe, P. P. et al. Decadal change in seabird-driven isotopes on islands with differing invasion histories. *Ecol. Appl.* **35**, e70030 (2025).
152. Miller-ter Kuile, A. et al. Impacts of rodent eradication on seed predation and plant community biomass on a tropical atoll. *Biotropica* **53**, 232–242 (2021).
153. Graham, N. A., Benkwitt, C. E. & Jones, H. P. Species eradication for ecosystem restoration. *Curr. Biol.* **34**, R407–R412 (2024).
154. Roberts, C. M., Duncan, R. P. & Wilson, K.-J. Burrowing seabirds affect forest regeneration, Rangitira Island, Chatham Islands, New Zealand. *N. Z. J. Ecol.* **31**, 208–222 (2007).
155. Berr, T. et al. Seabird and reef conservation must include coral islands. *Trends Ecol. Evol.* **38**, 490–494 (2023).
156. Steibl, S. et al. Atolls are globally important sites for tropical seabirds. *Nat. Ecol. Evol.* <https://doi.org/10.1038/s41559-024-02496-4> (2024).

157. Beetham, E., Kench, P. S. & Popinet, S. Future reef growth can mitigate physical impacts of sea-level rise on atoll islands. *Earth's Future* **5**, 1002–1014 (2017).
158. Steibl, S. et al. Rethinking atoll futures: local resilience to global challenges. *Trends Ecol. Evol.* **39**, 258–266 (2024).
159. Kench, P. S., Brander, R. W., Parnell, K. E. & McLean, R. F. Wave energy gradients across a Maldivian atoll: implications for island geomorphology. *Geomorphology* **81**, 1–17 (2006).
160. Toth, L. T. et al. The potential for coral reef restoration to mitigate coastal flooding as sea levels rise. *Nat. Commun.* **14**, 2313 (2023).
161. Hughes, T. P. et al. Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science* **359**, 80–83 (2018).
162. Kappes, P. J. et al. Do invasive mammal eradications from islands support climate change adaptation and mitigation? *Climate* **9**, 172 (2021).
163. Honzák, M. et al. Toward the quantification of the climate co-benefits of invasive mammal eradication on islands: a scalable framework for restoration monitoring. *Environ. Res. Lett.* **19**, 114018 (2024).
164. Longley-Wood, K., Engels, M., Lafferty, K. D., McLaughlin, J. P. & Wegmann, A. Transforming Palmyra Atoll to native-tree dominance will increase net carbon storage and reduce dissolved organic carbon reef runoff. *PLoS ONE* **17**, e0262621 (2022).
165. Oppel, S. et al. Estimating population size of a nocturnal burrow-nesting seabird using acoustic monitoring and habitat mapping. *Nat. Conserv.* **7**, 1–13 (2014).
166. Borker, A. L. et al. Do soundscape indices predict landscape-scale restoration outcomes? A comparative study of restored seabird island soundscapes. *Restor. Ecol.* **28**, 252–260 (2020).
167. Harris, S. A., Shears, N. T. & Radford, C. A. Ecoacoustic indices as proxies for biodiversity on temperate reefs. *Methods Ecol. Evol.* **7**, 713–724 (2016).
168. Dunn, R. E. Atolls are vital for seabirds and vice versa. *Nat. Ecol. Evol.* **8**, 1784–1785 (2024).
169. Mizutani, H., Kabaya, Y., Moors, P. J., Speir, T. W. & Lyon, G. L. Nitrogen isotope ratios identify deserted seabird colonies. *Auk* **108**, 960–964 (1991).
170. Pascoe, P., Shaw, J., Trebilco, R., Kong, S. & Jones, H. Island characteristics and sampling methodologies influence the use of stable isotopes as an ecosystem function assessment tool. *Ecol. Solut. Evid.* **2**, e12082 (2021).
171. Sato, N. et al. The distinctive material cycle associated with seabirds and land crabs on a pristine oceanic island: a case study of Minamiwoto, Ogasawara Islands, subtropical Japan. *Oecologia* **207**, 88 (2025).
172. Pascoe, P. et al. Temporal and spatial variability in stable isotope values on seabird Islands: what, where and when to sample. *Ecol. Indic.* **143**, 109344 (2022).
173. Stergiou, K. I. & Browman, H. I. Imbalances in the reporting and teaching of ecology from limnetic, oceanic and terrestrial eco-domains. *Mar. Ecol. Prog. Ser.* **304**, 292–297 (2005).
174. Rotjan, R. D. & Idjadi, J. Surf and turf: toward better synthesis by cross-system understanding. *Oikos* **122**, 285–287 (2013).
175. Di Marco, M. et al. Changing trends and persisting biases in three decades of conservation science. *Glob. Ecol. Conserv.* **10**, 32–42 (2017).
176. Morten, J. M. et al. Global marine flyways identified for long-distance migrating seabirds from tracking data. *Glob. Ecol. Biogeogr.* **34**, e70004 (2025).
177. Roman, J. et al. Migrating baleen whales transport high-latitude nutrients to tropical and subtropical ecosystems. *Nat. Commun.* **16**, 2125 (2025).
178. Jauharee, A. R. & Adam, M. S. *Significance of Seabirds to the Maldivian Tuna Fishery* (Indian Ocean Tuna Commission, 2012).
179. Signa, G., Mazzola, A. & Vizzini, S. Seabird influence on ecological processes in coastal marine ecosystems: an overlooked role? A critical review. *Estuar. Coast. Shelf Sci.* **250**, 107164 (2021).
180. Tobias, J. A. et al. AVONET: morphological, ecological and geographical data for all birds. *Ecol. Lett.* **25**, 581–597 (2022).
181. Carneiro, A. P. et al. The BirdLife Seabird Tracking Database: 20 years of collaboration for marine conservation. *Biol. Conserv.* **299**, 110813 (2024).
182. Young, H. S., Miller-ter Kuile, A., McCauley, D. J. & Dirzo, R. Cascading community and ecosystem consequences of introduced coconut palms (*Cocos nucifera*) in tropical islands. *Can. J. Zool.* **95**, 139–148 (2017).
183. Jones, H. P. et al. Severity of the effects of invasive rats on seabirds: a global review. *Conserv. Biol.* **22**, 16–26 (2008).
184. Towns, D. et al. in *Seabird Islands: Ecology, Invasion, and Restoration* (eds Mulder, C. P. H. et al.) 56–90 (Oxford Univ. Press, 2011).
185. Spatz, D. R. et al. in *Conservation of Marine Birds* (eds Young, L. & VanderWerf, E.) 97–130 (Academic Press, 2023).
186. Lorrain, A. et al. Seabirds supply nitrogen to reef-building corals on remote pacific islets. *Sci. Rep.* **7**, 3721 (2017).
187. Choinsard, N. et al. Tracing the fate of seabird-derived nitrogen in a coral reef using nitrate and coral skeleton nitrogen isotopes. *Limnol. Oceanogr.* **69**, 309–324 (2024).
188. Delevaux, J. M. et al. A linked land–sea modeling framework to inform ridge-to-reef management in high oceanic islands. *PLoS ONE* **13**, e0193230 (2018).
189. Thorne, L. H., Clay, T. A., Phillips, R. A., Silvers, L. G. & Wakefield, E. D. Effects of wind on the movement, behavior, energetics, and life history of seabirds. *Mar. Ecol. Prog. Ser.* **723**, 73–117 (2023).
190. Shepard, E., Cole, E.-L., Neate, A., Lempidakis, E. & Ross, A. Wind prevents cliff-breeding birds from accessing nests through loss of flight control. *eLife* **8**, e43842 (2019).
191. Leichter, J. J., Stewart, H. L. & Miller, S. L. Episodic nutrient transport to Florida coral reefs. *Limnol. Oceanogr.* **48**, 1394–1407 (2003).
192. Williams, G. J. et al. Biophysical drivers of coral trophic depth zonation. *Mar. Biol.* **165**, 60 (2018).
193. Steven, A. D. L. & Atkinson, M. J. Nutrient uptake by coral-reef microatolls. *Coral Reefs* **22**, 197–204 (2003).
194. Dunn, R. E. et al. Island restoration to rebuild seabird populations and amplify coral reef functioning. *Conserv. Biol.* **39**, e14313 (2025).
195. Stuart, C. et al. Seascape configuration determines spatial patterns of seabird-vectored nutrient enrichment to coral reefs. *Ecography* **2025**, e07863 (2025).
196. Morais, R. A., Patricio-Valerio, L., Narvaez, P., Parravicini, V. & Brandl, S. J. Rethinking Darwin's coral reef paradox and the ubiquity of 'marine oases'. *Curr. Biol.* **35**, 3241–3250.e6 (2025).
197. Polis, G. A., Hurd, S. D., Jackson, C. T. & Piñero, F. S. El Niño effects on the dynamics and control of an island ecosystem in the Gulf of California. *Ecology* **78**, 1884–1897 (1997).
198. Schmidt, S., Dennison, W. C., Moss, G. J. & Stewart, G. R. Nitrogen ecophysiology of Heron Island, a subtropical coral cay of the great barrier reef, Australia. *Funct. Plant. Biol.* **31**, 517–528 (2004).
199. Tomlinson, S. et al. Applications and implications of ecological energetics. *Trends Ecol. Evol.* **29**, 280–290 (2014).
200. Malhi, Y. et al. Logged tropical forests have amplified and diverse ecosystem energetics. *Nature* **612**, 707–713 (2022).
201. Edney, A. J. & Wood, M. J. Applications of digital imaging and analysis in seabird monitoring and research. *Ibis* **163**, 317–337 (2021).
202. Gauthreaux, S. A. Jr. & Belser, C. G. Radar ornithology and biological conservation. *Auk* **120**, 266–277 (2003).
203. Goddijn-Murphy, L., O'Hanlon, N. J., James, N. A., Masden, E. A. & Bond, A. L. Earth observation data for seabirds and their habitats: an introduction. *Remote Sens. Appl. Soc. Environ.* **24**, 100619 (2021).
204. Fretwell, P. T. & Trathan, P. N. Discovery of new colonies by Sentinel2 reveals good and bad news for emperor penguins. *Remote Sens. Ecol. Conserv.* **7**, 139–153 (2021).
205. Brodie, J., De'ath, G., Devlin, M., Furnas, M. & Wright, M. Spatial and temporal patterns of near-surface chlorophyll a in the Great Barrier Reef lagoon. *Mar. Freshw. Res.* **58**, 342–353 (2007).
206. Suryan, R. M., Santora, J. A. & Sydeman, W. J. New approach for using remotely sensed chlorophyll a to identify seabird hotspots. *Mar. Ecol. Prog. Ser.* **451**, 213–225 (2012).
207. Rakotoarivony, M. N. A. et al. Using imaging spectroscopy to assess the impacts of invasive plants on aboveground and belowground characteristics. *GIScience Remote Sens.* **61**, 2399388 (2024).
208. Smith, M.-L. et al. Direct estimation of aboveground forest productivity through hyperspectral remote sensing of canopy nitrogen. *Ecol. Appl.* **12**, 1286–1302 (2002).
209. Thomson, E. R. et al. Multiscale mapping of plant functional groups and plant traits in the high arctic using field spectroscopy, UAV imagery and Sentinel-2A data. *Environ. Res. Lett.* **16**, 055006 (2021).
210. Wang, Z. et al. Vegetation indices for mapping canopy foliar nitrogen in a mixed temperate forest. *Remote Sens.* **8**, 491 (2016).
211. Noppen, L. et al. Constraining industrial ammonia emissions using hyperspectral infrared imaging. *Remote Sens. Environ.* **291**, 113559 (2023).
212. Garcia-Quintas, A. et al. Machine and deep learning approaches to understand and predict habitat suitability for seabird breeding. *Ecol. Evol.* **13**, e10549 (2023).
213. Chapuis, L., Williams, B., Gordon, T. A. C. & Simpson, S. D. Low-cost action cameras offer potential for widespread acoustic monitoring of marine ecosystems. *Ecol. Indic.* **129**, 107957 (2021).
214. Williams, B. et al. Enhancing automated analysis of marine soundscapes using ecoacoustic indices and machine learning. *Ecol. Indic.* **140**, 108986 (2022).
215. Borker, A. L. *Applying Ecoacoustics to Bird Conservation and Monitoring* (UC Santa Cruz, 2018).
216. Penar, W., Magiera, A. & Kłoczek, C. Applications of bioacoustics in animal ecology. *Ecol. Complex.* **43**, 100847 (2020).
217. Lamont, T. A. C. et al. The sound of recovery: coral reef restoration success is detectable in the soundscape. *J. Appl. Ecol.* **59**, 742–756 (2022).
218. McInnes, J. C., Bird, J. P., Deagle, B. E., Polanowski, A. M. & Shaw, J. D. Using DNA metabarcoding to detect burrowing seabirds in a remote landscape. *Conserv. Sci. Pract.* **3**, e439 (2021).
219. de Leeuw, J. J., van den Brink, X., Gabrielsen, G. W. & Nijland, R. DNA metabarcoding reveals high diversity of fish and macrofaunal species in diets of little auks and other Arctic seabird species in Svalbard. *Polar Reefs* **47**, 1013–1023 (2024).
220. Duda, M. P. et al. Reconstructing long-term changes in avian populations using lake sediments: opening a window onto the past. *Front. Ecol. Evol.* **9**, 698175 (2021).
221. Davis, R. A., Seddon, P. J., Craig, M. D. & Russell, J. C. A review of methods for detecting rats at low densities, with implications for surveillance. *Biol. Invasions* **25**, 3773–3791 (2023).
222. Piaggio, A. J. et al. Evaluation of environmental DNA as a surveillance tool for invasive house mice (*Mus musculus*). *Environ. DNA* **7**, e70069 (2025).
223. Clay, C. G. et al. Exploring species and functional diversity of fishes in Cambodian coastal habitats using eDNA metabarcoding. *Coral Reefs* **44**, 221–241 (2025).
224. Gallego, R., Jacobs-Palmer, E., Cribari, K. & Kelly, R. P. Environmental DNA metabarcoding reveals winners and losers of global change in coastal waters. *Proc. R. Soc. B Biol. Sci.* **287**, 20202424 (2020).
225. Wright, D. G., van der Wal, R., Wanless, S. & Bardgett, R. D. The influence of seabird nutrient enrichment and grazing on the structure and function of island soil food webs. *Soil. Biol. Biochem.* **42**, 592–600 (2010).

226. Glasl, B., Webster, N. S. & Bourne, D. G. Microbial indicators as a diagnostic tool for assessing water quality and climate stress in coral reef ecosystems. *Mar. Biol.* **164**, 91 (2017).
227. Cordier, T. et al. Ecosystems monitoring powered by environmental genomics: a review of current strategies with an implementation roadmap. *Mol. Ecol.* **30**, 2937–2958 (2021).

Acknowledgements

The authors thank the David and Lucile Packard Foundation (D.J.W.), the Bertarelli Foundation as part of the Bertarelli Programme in Marine Science (C.E.B., N.A.J.G. and Y.M.), the Alexander von Humboldt Foundation (S.S.), the Leverhulme Trust (Y.M.) and the NERC ENVISION doctoral training programme (L.-L.J.) for funding.

Author contributions

H.P.J. and N.A.J.G. amassed the co-authors of the article. H.P.J. led initial discussions of the article structure and content. The authors contributed equally to all other aspects of the article.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s44358-025-00099-w>.

Peer review information *Nature Reviews Biodiversity* thanks Leandro Bugoni and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© Springer Nature Limited 2025